


In the Claims:

1. **(previously presented)** A compound represented by formula I:



I

wherein,

n is 1, 3, or 4;

R represents independently for each occurrence H, alkyl, aryl, -CH₂-aryl, -C(O)-alkyl, -C(O)-aryl, or -Si(alkyl)₃;

R¹ and R² are independently H, -CH₂-aryl, -C(O)-alkyl, -C(O)-aryl, -Si(alkyl)₃; or R¹ and R² taken together are C(CH₃)₂, P(O)OH, or P(O)OR⁵;

R³ is amino, -N₃, or -NH₃X;


R⁴ represents independently for each occurrence H, alkyl, aryl, -CH₂-aryl, -C(O)-alkyl, -C(O)-aryl, -Si(alkyl)₃, or -P(O)(OR⁵)₂;


R⁵ represents independently for each occurrence H, Li⁺, Li⁺, Na⁺, K⁺, Rb⁺, Cs⁺, aryl, or an optionally substituted alkyl group; and

X is a halogen, alkyl carboxylate, or aryl carboxylate.


2. **(canceled)**
3. **(original)** The compound of claim 1, wherein n is 3.
4. **(original)** The compound of claim 1, wherein R is H.

5. **(original)** The compound of claim 1, wherein R^1 and R^2 taken together are $P(O)OR^5$.
6. **(original)** The compound of claim 1, wherein R^3 is N_3 .
7. **(original)** The compound of claim 1, wherein R^3 is $-NH_3X$.
8. **(original)** The compound of claim 1, wherein R^4 represents independently for each occurrence H, $-CH_2Ph$, or $-Si(alkyl)_3$.
9. **(original)** The compound of claim 1, wherein R^4 represents independently for each occurrence H, $-CH_2Ph$, -or $P(O)OR^5$; and R^5 is an optionally substituted alkyl group.
10. **(currently amended)** A compound selected from the group consisting of:







5




5




5






5







5







5
























11. **(previously presented)** A compound represented by formula **II**:



II

wherein,



n is 1, 3, or 4;


R represents independently for each occurrence H, alkyl, aryl, -CH₂-aryl, -C(O)-alkyl, -C(O)-aryl, or -Si(alkyl)₃;

R¹ is -(CH₂)_mCH=CH₂ or trichloroacetimidate; and

m is 1-6.

12. **(canceled)**
13. **(original)** The compound of claim 11, wherein n is 3.
14. **(original)** The compound of claim 11, wherein m is 3.
15. **(original)** The compound of claim 11, wherein R represents independently for each occurrence -CH₂-aryl or -Si(alkyl)₃.
16. **(original)** The compound of claim 11, wherein R represents independently for each occurrence benzyl or -Si(iPr)₃.
17. **(currently amended)** The compound of claim 11, wherein R¹ is trichloroacetimidate and R represents independently for each occurrence benzyl or -Si(iPr)₃. ~~and~~
18. **(currently amended)** The compound of claim 11, wherein said compound of formula **II** is selected from the group consisting of:





- R7O[C@H]1[C@H](OR)[C@H](O[C@@H]1SR6)R3

[[,]] with a

A chemical structure of a branched oligosaccharide. It consists of three pyranose rings. The top ring is a 6-membered ring with an OR group at the C6 position. It is linked via an oxygen atom to the C1 position of a middle 6-membered ring. The middle ring has an OR group at the C2 position and is linked via an oxygen atom to the C1 position of a bottom 6-membered ring. The bottom ring has OR groups at the C2 and C3 positions. Various other OR groups are attached to the rings, labeled R¹O, R²O, R³O, R⁷O, and RO.

$$\text{R}^2\text{O}-\text{C}_4\text{H}_8-\text{OR}^1; \text{ wherein,}$$

R¹ and R² are independently H, -CH₂-aryl, -C(O)-alkyl, -C(O)-aryl, -Si(alkyl)₃; or R¹ and R² taken together are C(CH₃)₂, P(O)OH, or P(O)OR⁵;

R³ is amino, -N₃, or -NH₃X;

R⁵ represents independently for each occurrence H, Li⁺, Li⁺, Na⁺, K⁺, Rb⁺, Cs⁺, aryl, or an optionally substituted alkyl group;

R⁶ is alkyl or aryl;

R⁷ is alkyl, aryl, -CH₂-aryl, -C(O)-alkyl, -C(O)-aryl, or -Si(alkyl)₃; and

X is a halogen, alkyl carboxylate, or aryl carboxylate.

20. **(original)** The method of claim 19, wherein R is -CH₂-aryl.
21. **(original)** The method of claim 19, wherein R¹ and R² taken together are C(CH₃)₂.
22. **(original)** The method of claim 19, wherein R³ is -N₃.
23. **(original)** The method of claim 19, wherein R⁶ is alkyl.
24. **(original)** The method of claim 19, wherein R⁷ is -C(O)-alkyl.
25. **(original)** The method of claim 19, wherein R is benzyl, R¹ and R² taken together are C(CH₃)₂, and R³ is -N₃.
26. **(original)** The method of claim 19, wherein R is benzyl, R¹ and R² taken together are C(CH₃)₂, R³ is -N₃, and R⁶ is ethyl.
27. **(currently amended)** A method of preparing a tetrasaccharide, comprising the steps of:
covalently binding a mannopyranoside to a solid support to provide a first substrate, reacting said first substrate with a mannopyranose trichloroacetimidate to give a disaccharide bound to said solid support, reacting said disaccharide with a mannopyranose trichloroacetimidate to give a trisaccharide bound to said solid support, reacting said trisaccharide with a mannopyranose trichloroacetimidate to give a tetrasaccharide bound to said solid support, and cleaving said tetrasaccharide from said solid support.
28. **(original)** The method of claim 27, wherein said mannopyranoside is bound to said solid support through a glycosidic linkage.
29. **(original)** The method of claim 27, wherein said tetrasaccharide is cleaved from said solid support using Grubbs' catalyst.

30. (currently amended) The method of claim 27, wherein said tetrasaccharide is

